

UC Davis

UC Davis Previously Published Works

Title

Draft Genome Sequence of Multidrug-Resistant Abortive *Campylobacter jejuni* from Northern California.

Permalink

<https://escholarship.org/uc/item/99g217m7>

Journal

Genome announcements, 5(15)

ISSN

2169-8287

Authors

Weis, Allison M
Clothier, Kristin A
Huang, Bihua C
et al.


Publication Date

2017-04-01

DOI

10.1128/genomea.00171-17

Peer reviewed



Draft Genome Sequence of Multidrug-Resistant Abortive *Campylobacter jejuni* from Northern California

Allison M. Weis,^a Kristin A. Clothier,^b Bihua C. Huang,^a Nguyet Kong,^a

 Bart C. Weimer^a

School of Veterinary Medicine, 100K Pathogen Genome Project, UC Davis, Davis, California, USA^a; California Animal Health and Food Safety, Davis, California, USA^b

ABSTRACT *Campylobacter jejuni* is an enteric bacterium that can cause abortion in livestock. This is the release of a multidrug-resistant *Campylobacter jejuni* genome from an isolate that caused an abortion in a cow in northern California. This isolate is part of the 100K Pathogen Genome Project.

Campylobacter jejuni is an enteric pathogen that affects humans worldwide with infection accompanied by symptoms such as fever and bloody diarrhea, and can lead to autoimmune diseases (1–6). *C. jejuni* is zoonotic, and in livestock it can lead to gastroenteritis and occasionally abortion (7–12). In recent years, several regions in the United States have noted the emergence of an abortive hypervirulent *C. jejuni* isolate (13–15).

Multidrug resistant bacteria are a major worldwide problem, leading to untreatable infection and mortality among humans and other animals (16, 17) that can be transmitted to the human microbiome (18). Here, we describe a multidrug-resistant abortive *C. jejuni* strain from northern California that was isolated from a bovine fetus from a third trimester abortion in 2009 and was uniquely multidrug resistant. This *C. jejuni* genome contains *tetO*, which confers resistance to tetracycline and its derivatives. It also contains *aphA*, a kanamycin resistance gene that is an aminoglycoside phosphotransferase. Lastly, it also contains a Thr-86-Ile mutation in the GyrA protein that results in fluoroquinolone resistance. The *tetO* and *aphA* genes are located on a putative plasmid, whereas *gyrA* is in the chromosome. The genome assembly comprised 54 contigs, 1,708,778 bp, and 1,717 coding sequences, and the sequence depth was 59×. This *C. jejuni* isolate was among a set of animal source isolates that was given to the 100K Pathogen Genome Project (<http://www.100kgenomes.org>) in the laboratory of Bart Weimer (UC Davis). The isolate was checked for purity (19) and genomic DNA (gDNA) was extracted from culture grown on 5% blood agar plates (UC Davis, VetMed Biological Services) for 1 to 2 days, lysed with an enzyme cocktail (20), purified with a Qiagen QIAamp DNA minikit (51306), and analyzed on the Agilent 2200 TapeStation system with the Genomic DNA ScreenTape assay for integrity of gDNA, as described previously (21–23). Approximately one microgram of gDNA was used for library construction with the KAPA Hyper library preparation kit (Kapa Biosystems, Boston, MA, USA; KK8514), on a PerkinElmer Sciclone NGS Workstation (Waltham, MA, USA). In this process, fragmented double-stranded gDNA molecules were end-repaired (5'), adenylated (3'), and ligated with double stranded DNA adapters (Weimer 384 TS-LT DNA Barcodes, Integrated DNA Technologies, Inc., Coralville, IA, USA), followed by dual-SPRI size selection for targeting a library size of 250 to 500 bp. Final library quality assessment to verify size distribution was performed on a Perkin Elmer Caliper Lab Chip GX using the PerkinElmer Caliper HT 1K kit (CLS701879), and library quantification was completed using a SYBR Green–based qPCR kit and a KAPA library quantification kit

Received 13 February 2017 **Accepted** 21 February 2017 **Published** 13 April 2017

Citation Weis AM, Clothier KA, Huang BC, Kong N, Weimer BC. 2017. Draft genome sequence of multidrug-resistant abortive *Campylobacter jejuni* from northern California. Genome Announc 5:e00171-17. <https://doi.org/10.1128/genomeA.00171-17>.

Copyright © 2017 Weis et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Bart C. Weimer, bcweimer@ucdavis.edu.

(KK4824). Pooled libraries were sequenced on the Illumina HiSeq X TEN platform with a PE150-plus index read at Novogene (Sacramento, CA, USA). Paired-end sequence reads were assembled using ABySS version 1.9.0 (k -mer = 64) and annotated with Prokka (24, 25).

Accession number(s). This sequence can be found at NCBI SRA BioProject PRJNA186441 under genome SRA accession number SRR5210995, and at NCBI GenBank under the accession number [MUJW00000000](#).

ACKNOWLEDGMENTS

We thank all contributors of the 100K Pathogen Genome Project and members of the Weimer Lab for technical assistance. We thank the individuals associated with the isolation of the unique and important *C. jejuni* strain presented here. This project was funded by the 100K Pathogen Genome Project to produce this sequence.

REFERENCES

- Coker AO, Isokpehi RD, Thomas BN, Amisu KO, Obi CL. 2002. Human campylobacteriosis in developing countries. *Emerg Infect Dis* 8:237–244. <https://doi.org/10.3201/eid0803.010233>.
- CDC. 2012. Foodborne diseases Active Surveillance Network (FoodNet): FoodNet surveillance report for 2011 (final report). Department of Health and Human Services, CDC, Atlanta, GA. https://www.cdc.gov/foodnet/pdfs/2011_annual_report_508c.pdf.
- Nachamkin I. 2003. *Campylobacter* and *Arcobacter*, p 907. In Murray E, Baron J, Jorgensen JH, Pfaller MA, Tenover FC, Tenover FC (ed), *Manual of clinical microbiology*, 8th ed. ASM Press, Washington, DC.
- Altekruse SF, Stern NJ, Fields PI, Swerdlow DL. 1999. *Campylobacter jejuni*—an emerging foodborne pathogen. *Emerg Infect Dis* 5:28–35. <https://doi.org/10.3201/eid0501.990104>.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, Griffin PM. 2011. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis* 17:7–15. <https://doi.org/10.3201/eid1701.P11101>.
- Young KT, Davis LM, DiRita VJ. 2007. *Campylobacter jejuni*: molecular biology and pathogenesis. *Nat Rev Microbiol* 5:665–679. <https://doi.org/10.1038/nrmicro1718>.
- Nielsen EM. 2002. Occurrence and strain diversity of thermophilic campylobacters in cattle of different age groups in dairy herds. *Lett Appl Microbiol* 35:85–89. <https://doi.org/10.1046/j.1472-765X.2002.01143.x>.
- Jacobs-Reitsma W, Lyhs U, Wagenaar J. 2008. *Campylobacter* in the food supply, p 627–644. In Nachamkin I, Szymanski CM, Blaser MJ (ed), *Campylobacter*, 3rd ed. ASM Press, Washington, DC.
- Weis AM, Miller WA, Byrne BA, Chouicha N, Boyce WM, Townsend AK. 2014. Prevalence and pathogenic potential of *Campylobacter* isolates from free-living, human-commensal American crows. *Appl Environ Microbiol* 80:1639–1644. <https://doi.org/10.1128/AEM.03393-13>.
- Taff CC, Weis AM, Wheeler S, Hinton MG, Weimer BC, Barker CM, Jones M, Logsdon R, Smith WA, Boyce WM, Townsend AK. 2016. Influence of host ecology and behavior on *Campylobacter jejuni* prevalence and environmental contamination risk in a synanthropic wild bird species. *Appl Environ Microbiol* 82:4811–4820. <https://doi.org/10.1128/AEM.01456-16>.
- Gardner TJ, Fitzgerald C, Xavier C, Klein R, Pruckler J, Stroika S, McLaughlin JB. 2011. Outbreak of campylobacteriosis associated with consumption of raw peas. *Clin Infect Dis* 53:26–32. <https://doi.org/10.1093/cid/cir249>.
- Weis AM, Storey DB, Taff CC, Townsend AK, Huang BC, Kong NT, Clothier KA, Spinner A, Byrne BA, Weimer BC. 2016. Genomic comparison of *Campylobacter* spp. and their potential for zoonotic transmission between birds, primates, and livestock. *Appl Environ Microbiol* 82:7165–7175. <https://doi.org/10.1128/AEM.01746-16>.
- Sahin O, Fitzgerald C, Stroika S, Zhao S, Sippy RJ, Kwan P, Plummer PJ, Han J, Yaeger MJ, Zhang Q. 2012. Molecular evidence for zoonotic transmission of an emergent, highly pathogenic *Campylobacter jejuni* clone in the United States. *J Clin Microbiol* 50:680–687. <https://doi.org/10.1128/JCM.06167-11>.
- Sahin O, Plummer PJ, Jordan DM, Sulaj K, Pereira S, Robbe-Austerman S, Wang L, Yaeger MJ, Hoffman LJ, Zhang Q. 2008. Emergence of a tetracycline-resistant *Campylobacter jejuni* clone associated with outbreaks of ovine abortion in the United States. *J Clin Microbiol* 46:1663–1671. <https://doi.org/10.1128/JCM.00031-08>.
- Hedstrom OR, Sonn RJ, Lassen ED, Hultgren BD, Crisman RO, Smith BB, Snyder SP. 1987. Pathology of *Campylobacter jejuni* abortion in sheep. *Vet Pathol* 24:419–426. <https://doi.org/10.1177/030098588702400509>.
- Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S, Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens MJ, Vatopoulos A, Weber JT, Monnet DL. 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 18:268–281. <https://doi.org/10.1111/j.1469-0691.2011.03570.x>.
- Nikaido H. 2009. Multidrug resistance in bacteria. *Annu Rev Biochem* 78:119–146. <https://doi.org/10.1146/annurev.biochem.78.082907.145923>.
- Hu Y, Yang X, Li J, Lv N, Liu F, Wu J, Lin IY, Wu N, Weimer BC, Gao GF, Liu Y, Zhu B. 2016. The bacterial mobile Resistome Transfer Network connecting the animal and human microbiomes. *Appl Environ Microbiol* 82:6672–6681. <https://doi.org/10.1128/AEM.01802-16>.
- Kong N, Ng W, Lee V, Kelly L, Weimer BC. 2013. Production and analysis of high molecular weight genomic DNA for NGS pipelines using Agilent DNA extraction kit (p/n 200600). Agilent Technologies Application Note. <https://doi.org/10.13140/RG.2.1.2961.4807>.
- Jeannotte R, Lee E, Kong N, Ng W, Kelly L, Weimer BC. 2014. High-throughput analysis of foodborne bacterial genomic DNA using Agilent 2200 TapeStation and genomic DNA ScreenTape system. Agilent Technologies Application Note. <https://doi.org/10.13140/RG.2.1.3354.6961>.
- Kong N, Ng W, Cai L, Leonardo A, Kelly L, Weimer BC. 2014. Integrating the DNA integrity number (DIN) to assess genomic DNA (gDNA) quality control using the Agilent 2200 TapeStation system. Agilent Technologies Application Note. <https://doi.org/10.13140/RG.2.1.3616.8409>.
- Weis AM, Huang BC, Storey DB, Kong N, Chen P, Arabyan N, Gilpin B, Mason C, Townsend AK, Smith WA, Byrne BA, Taff CC, Weimer BC. 2017. Large-scale release of campylobacter draft genomes: resources for food safety and public health from the 100K pathogen genome project. *Genome Announc* 5(1):e00925-16. <https://doi.org/10.1128/genomeA.00925-16>.
- Weis AM, Clothier KA, Huang BC, Kong N, Weimer BC. 2016. Draft genome sequences of *Campylobacter jejuni* strains that cause abortion in livestock. *Genome Announc* 4(6):e01324-16. <https://doi.org/10.1128/genomeA.01324-16>.
- Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJM, Birol I. 2009. ABySS: a parallel assembler for short read sequence data. *Genome Res* 19:1117–1123. <https://doi.org/10.1101/gr.089532.108>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.